

Riemann surfaces and AF -algebras

Igor Nikolaev *

Abstract

For a generic set in the Teichmüller space, we construct a covariant functor with the range in a category of the AF -algebras. The functor maps isomorphic Riemann surfaces to the stably isomorphic AF -algebras. Our construction is based on a Hodge theory for measured foliations, elaborated by Hubbard, Masur and Thurston. As a special case, one obtains a categorical correspondence between the complex and noncommutative tori.

Key words and phrases: Riemann surfaces, AF -algebras

AMS (MOS) Subj. Class.: 14H10, 46L40, 58F10

1 Introduction

1. Setting of problems.

A. Let $\mathbb{C}$ be the complex plane and $\tau \in \mathbb{C}$ a point, such that $\text{Im } \tau > 0$. Denote by $\mathcal{B}$ a category whose objects, $\text{Ob } (\mathcal{B})$, are the complex tori $\mathbb{B}_\tau = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ and morphisms, $\text{Mor } (\mathcal{B})$, are the biholomorphic maps modulo the ones, which are isotopic to the identity map with respect to a fixed topological marking. Likewise, let $\mathcal{A}$ be a category, whose objects $\text{Ob } (\mathcal{A})$ are the AF -algebras $\mathbb{A}_\theta$ (the *Effros-Shen* algebras [7]), given by the following Bratteli diagram:




Figure 1: The Effros-Shen algebra $\mathbb{A}_\theta$.

and morphisms $\text{Mor } (\mathcal{A})$ are the stable isomorphisms of the AF -algebras modulo the inner automorphisms. (Here θ is an irrational number, $a_0 \in \mathbb{N} \cup \{0\}$ and

*Partially supported by NSERC.

$a_i \in \mathbb{N}$ are the multiplicities of edges of the graph.) The following phenomenon is well known.

Observation. *For a generic $\mathbb{A}_\theta \in \mathcal{A}$ and $\mathbb{B}_\tau \in \mathcal{B}$, it holds $\text{Mor}(\mathbb{A}_\theta) \cong \text{Mor}(\mathbb{B}_\tau) \cong SL_2(\mathbb{Z})/\pm I$.*

(*Proof.* Indeed, $\mathbb{B}_\tau, \mathbb{B}_{\tau'}$ are isomorphic, if and only if, $\tau' = \frac{a\tau+b}{c\tau+d}$, where $ad-bc = 1$ and $a, b, c, d \in \mathbb{Z}$. Generically, the group of the inner automorphisms of $\mathbb{B}_\tau$ is $\pm I$, where I is the identity matrix. Similarly, the AF -algebras $\mathbb{A}_\theta, \mathbb{A}_{\theta'}$ are stably isomorphic, i.e. $\mathbb{A}_\theta \otimes \mathcal{K} \cong \mathbb{A}_{\theta'} \otimes \mathcal{K}$, where $\mathcal{K}$ is the C^* -algebra of compact operators, if and only if, $\theta' = \frac{a\theta+b}{c\theta+d}$, where $ad-bc = 1$ and $a, b, c, d \in \mathbb{Z}$. Again, for a generic $\mathbb{A}_\theta$ the group of stable inner automorphisms is $\pm I$. $\square$)

B. The paper is about this phenomenon and its higher genus manifestations. Our goal is a general correspondence between the Riemann surfaces and the AF -algebras. It is assumed at the outset, that the observed phenomenon is part of a categorical correspondence. Namely, recall that covariant functor is a pair of functions $F_1 : \text{Ob}(\mathcal{B}) \rightarrow \text{Ob}(\mathcal{A})$ and $F_2 : \text{Mor}(\mathcal{B}) \rightarrow \text{Mor}(\mathcal{A})$, denoted by the same symbol F , such that $F(\varphi_1\varphi_2) = F(\varphi_1)F(\varphi_2)$ for any $\varphi_1, \varphi_2 \in \text{Mor}(\mathcal{B})$. Our objective can be expressed as follows.

Main problem. *To construct a covariant functor $F : \mathcal{B} \rightarrow \mathcal{A}$ (if any), which maps isomorphic complex tori to the stably isomorphic Effros-Shen algebras.*

One can generalize the main problem as follows. Let $\mathcal{S}$ be a category whose objects, $\text{Ob}(\mathcal{S})$, are the Riemann surfaces $\mathbb{S}$ and morphisms, $\text{Mor}(\mathcal{S})$, are isomorphisms of the Riemann surfaces. Likewise, let $\mathcal{A}^*$ be a category whose objects, $\text{Ob}(\mathcal{A}^*)$, are certain AF -algebras, $\mathbb{A}$, which we shall call the *toric AF -algebras*, and morphisms, $\text{Mor}(\mathcal{A}^*)$, are stable isomorphisms of the AF -algebras.

Extended main problem. *Construct a covariant functor $F^* : \mathcal{S} \rightarrow \mathcal{A}^*$ (if any), which maps isomorphic Riemann surfaces to the stably isomorphic toric AF -algebras.*

2. Background.

A correspondence between the categories $\mathcal{A}$ and $\mathcal{B}$ can be traced back to F. Klein [12]. In a modern context, it was reported by Connes-Douglas-Schwarz [3], Kontsevich [13], Manin [15] and others. In the cited works, the $\mathcal{A}$ is viewed as a “noncommutative boundary” of $\mathcal{B}$, to which the action of the modular group extends. The philosophy $\mathcal{A} \cong \partial\mathcal{B}$ prevails in the works of Kontsevich-Soibelman [14], Polishchuk-Schwarz [19], Soibelman-Vologodsky [20] and others. Note that the authors prefer to work with the noncommutative tori, to which the Effros-Shen algebras are isomorphic from the standpoint of K -theory.

3. Objectives.

The extended functor $F^* : \mathcal{S} \rightarrow \mathcal{A}^*$ attracted less attention so far. The objective of present paper is detailed construction of such a functor. We build a

functor F^* , such that the restriction of F^* to the Riemann surfaces of genus one coincides with functor $F : \mathcal{B} \rightarrow \mathcal{A}$. Thus, we solve the main and the extended main problems. However, a difference between the two cases exists. Unless $g = 1$, functor F^* is defined only for a generic set in the Teichmüller space $T(g)$. This effect is due to a (non-generic) divergence of the Jacobi-Perron continued fractions. Let us note, that functor F^* can be extended to the non-generic cases as well. However, the group $Mor(\mathbb{A})$, $\mathbb{A} \in \mathcal{A}^*$ is no longer isomorphic to the matrix group $GL_{6g-6}(\mathbb{Z})$.

4. Construction of the mapping F^* .

A. Let us outline the construction of F^* . It is based on a Hodge theory for the measured foliations developed by Hubbard-Masur [11] and inspired by Thurston [22]. The measured foliation, $\mathcal{F}$, can be defined as a local-global structure on the surface X , consisting of a finite collection of the singular points x_i and regular leaves (1-dimensional submanifolds). Each x_i has a local chart V , such that $\mathcal{F} = \mathcal{F}(x_i, \phi_i)$ is given by the trajectories of a closed 1-form $\phi_i = Im(z^{\frac{k_i}{2}} dz)$, where $z = u + iv \in V$ and $k_i \in \mathbb{N}$. Everywhere else, $\mathcal{F}$ looks as a family of the parallel lines. It was observed by Jenkins and Strebel, that every measured foliation occurs uniquely as foliation $Im q = 0$ given by a holomorphic quadratic form q on the Riemann surface $\mathbb{S}$. This fact has been established for the foliations, whose regular leaves are homeomorphic to S^1 (the Jenkins-Strebel foliations, or JS -foliations), see e.g. [21]. The equivalence classes of the JS -foliations is a non-empty set of the real dimension $6g - 6$. The uniqueness of q implies (via a standard argument, see below), that the JS -foliations parametrize the Teichmüller space of surface X . The parametrization has been rapidly adopted by Harer, Kontsevich and Mumford to represent the Teichmüller space via the metric ribbon graphs, see e.g. §6 of [10] for an account.

B. Unlike the JS -foliation, each regular leaf of a minimal foliation is dense in X (by definition). Hubbard and Masur extended Jenkins-Strebel's results to the minimal foliations, which we shall call the Hubbard-Masur, or HM -foliations. A key step in Hubbard-Masur's proof involves the uniqueness of the JS -foliations, since they are dense in the space of all measured foliations. It is helpful to think of the HM -foliations as of the set of irrational numbers, while the JS -foliations form a set of the rational numbers of the real line. These two families of measured foliations are complementary to each other: the JS -foliations are topologically trivial, but rich in measure, while the HM -foliations are poor in measure, yet topologically rich. Let us give some details of the Hubbard-Masur construction.

C. Denote by $T(g)$ the Teichmüller space of genus $g \geq 1$ and let $\mathbb{S} \in T(g)$. Let $q \in H^0(\mathbb{S}, \Omega^{\otimes 2})$ be a holomorphic quadratic differential on the Riemann surface $\mathbb{S}$, such that all zeroes of q are simple. By $\tilde{\mathbb{S}}$ we understand a double cover of $\mathbb{S}$ ramified over the zeroes of q . Note that there is an involution on the homology groups $H_*(\tilde{\mathbb{S}})$ induced by the covering map $\tilde{\mathbb{S}} \rightarrow \mathbb{S}$. Let $H_1^{odd}(\tilde{\mathbb{S}})$ be the odd part of the first (integral) homology of $\tilde{\mathbb{S}}$ with respect to this involution, relatively

the zeroes of q . By the formulas for the relative homology:

$$H_1^{odd}(\tilde{\mathbb{S}}) \cong \mathbb{Z}^n, \text{ where } n = \begin{cases} 6g - 6, & \text{if } g \geq 2 \\ 2, & \text{if } g = 1. \end{cases}$$

D. Let $\mathcal{F}$ be a measured foliation with the simple zeroes (i.e. $k_i = 1$), Q the vector bundle of all quadratic differentials over the Teichmüller space $T(g)$ and 0 the zero section of Q . Consider those quadratic differentials, $E_{\mathcal{F}} \subset Q - \{0\}$, whose horizontal trajectories induce $\mathcal{F}$. Note that, if $\mathcal{F}$ is a measured foliation with the simple zeroes (a generic case), then $E_{\mathcal{F}} \cong \mathbb{R}^n - 0$, while $T(g) \cong \mathbb{R}^n$.

Hubbard-Masur homeomorphism ([11]). *There exists a homeomorphism (an embedding) $h : E_{\mathcal{F}} \rightarrow T(g)$.*

The Hubbard-Masur result implies that the measured foliations parametrize the space $T(g) - \{pt\}$, where $pt = h(0)$. Indeed, denote by $\mathcal{F}'$ a vertical trajectory structure $Re\ q = 0$ of q . Since $\mathcal{F}$ and $\mathcal{F}'$ define q , and $\mathcal{F} = Const$ for all $q \in E_{\mathcal{F}}$, one gets a homeomorphism between $T(g) - \{pt\}$ and Φ_X , where $\Phi_X \cong \mathbb{R}^n - 0$ is the space of equivalence classes of the measured foliations $\mathcal{F}'$ on X .

E. Recall that $T(g) \cong \mathbb{R}^n$, where $n = \dim H_1^{odd}(\tilde{\mathbb{S}})$. In view of the Hubbard-Masur homeomorphism, any n -parameter family of the measured foliations can be used to parametrize the space $T(g) - \{pt\}$. As explained, there are two notable families of such foliations: the *JS* and *HM*-foliations. Note that the equivalence classes of the *JS*-foliations form a dense countable subset in the space of all measured foliations. For the case of $g = 1$, the *JS*-foliations are given by the straight lines of a rational slope, parametrized by $\mathbb{Q}$. To better understand a difference between the *JS* and *HM*-foliations, let us denote them by $\mathcal{F}_{JS}$ and $\mathcal{F}_{HM}$, respectively. A *leaf space* of the foliation $\mathcal{F}$ on surface X is the factor space $X/\mathcal{F}$, which identifies the points of X belonging to the same leaf of $\mathcal{F}$. It is well known that generically (i.e. when all the zeroes of the foliation are simple), the space $X/\mathcal{F}_{JS} \cong \Gamma_n$, where Γ_n is a 3-valent metric graph embedded in X (a metric ribbon graph), see e.g. [10]. The edges of the graph correspond to the heights of the cylinders carved by the periodic trajectories of the *JS*-foliations, while the vertices are bijective with the zeroes of the foliation. An n -parameter family of the *JS*-foliations is obtained by a variation of the lengths of the edges of Γ_n . The graphs Γ_n can be used to calculate homology of the mapping class group (Harer) and to settle the Witten conjecture (Kontsevich).

F. Unlike the *JS*-foliations, the leaf space $X/\mathcal{F}_{HM}$ is no longer a 1-dimensional *CW*-complex (a graph), but a non-Hausdorff topological space. Due in part to this fact, the *HM*-foliations have been reluctantly used as a model for the Teichmüller spaces. As we shall see, the *AF*-algebras appear to be a proper category in this sense. Indeed, let $\lambda_i = \int_{\gamma_i} \phi$, where $\phi = Re\ q'$ is a closed 1-form and γ_i are elements of a basis in $H_1^{odd}(\tilde{\mathbb{S}})$. (The periods λ_i are well defined in the sense that a $\mathbb{Z}$ -module $\Lambda_n = \sum_{i=1}^n \lambda_i \mathbb{Z}$ depends neither on γ_i nor the equivalence class of the 1-form ϕ , see the proof of lemma 1.) It was shown by Douady and Hubbard in [4], that λ_i are coordinates in the space of all measured foliations.

Every n -parameter family of the HM -foliations is obtained by variation of the reals λ_i in a generic set of $\mathbb{R}^n$. Thus, λ_i form a coordinate system in a generic part of $T(g)$.

G. Let $Mod(X) = Diff^+(X) / Diff^0(X)$ be the mapping class group of the surface X . (For the case $g = 1$, $Mod(X) \cong SL_2(\mathbb{Z})$ is the familiar modular group, hence the notation.) The following $\mathbb{Z}$ -module, attached to an HM -foliation, will be critical. Assuming that λ_i are linearly independent over $\mathbb{Q}$, consider a $\mathbb{Z}$ -module:

$$\Lambda_n = \lambda_1 \mathbb{Z} + \dots + \lambda_n \mathbb{Z} \subset \mathbb{R},$$

thought of as a dense subset of the real line. One can ask what happens to $\Lambda_n = \Lambda_n(\mathbb{S})$, when one passes to an isomorphic Riemann surface $\mathbb{S}'$, i.e. $\mathbb{S}' = \varphi(\mathbb{S})$ for an $\varphi \in Mod(X)$? It is not hard to prove, that $\Lambda_n(\mathbb{S}) = \Lambda_n(\varphi(\mathbb{S}))$ for $\forall \varphi \in Mod(X)$, see lemma 1. (The action of φ corresponds to a change of basis in the module Λ_n .) In other words, the modules Λ_n make a category, which is covariant to the category $\mathcal{S}$. Roughly speaking, the rest of the text is a phrasing of this property of Λ_n in terms of the toric AF -algebras. Note that it is in a sharp contrast to the ribbon graph picture, where $\Gamma_n(\mathbb{S}) \neq \Gamma_n(\varphi(\mathbb{S}))$ for almost all $\varphi \in Mod(X)$.

H. To finalize construction of the mapping $F^* : \mathcal{S} \rightarrow \mathcal{A}^*$, let us denote by $\theta = (\theta_1, \dots, \theta_{n-1})$ a vector with the coordinates $\theta_i = \lambda_{i-1}/\lambda_1$, whenever $\lambda_1 \neq 0$. (Note that $\lambda_1 \neq 0$ is a generic restriction.) Consider the Jacobi-Perron continued fraction:

$$\begin{pmatrix} 1 \\ \theta \end{pmatrix} = \lim_{k \rightarrow \infty} \begin{pmatrix} 0 & 1 \\ I & b_1 \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ I & b_k \end{pmatrix} \begin{pmatrix} 0 \\ \mathbb{I} \end{pmatrix},$$

where $b_i = (b_1^{(i)}, \dots, b_n^{(i)})^T$ is a vector of the non-negative integers, I the unit matrix and $\mathbb{I} = (0, \dots, 0, 1)^T$. Denote by $\mathbb{A} := \mathbb{A}_{\theta_1, \dots, \theta_{n-1}}$ an AF -algebra:

$$\mathbb{Z}^n \xrightarrow{\begin{pmatrix} 0 & 1 \\ I & b_1 \end{pmatrix}} \mathbb{Z}^n \xrightarrow{\begin{pmatrix} 0 & 1 \\ I & b_2 \end{pmatrix}} \mathbb{Z}^n \dots,$$

which is defined as the inductive limit of the positive isomorphisms given by the unimodular integral matrices $\begin{pmatrix} 0 & 1 \\ I & b_i \end{pmatrix}$ for $i = 1, \dots, \infty$. The map F^* is given by the formula $\mathbb{S}(\lambda_1, \dots, \lambda_n) \mapsto \mathbb{A}_{\theta_1, \dots, \theta_{n-1}}$, where $\mathbb{S} \in \mathcal{S}$ and $\mathbb{A} \in \mathcal{A}$. Note that in the case $g = 1$, the AF -algebra $\mathbb{A}$ coincides with an Effros-Shen algebra $\mathbb{A}_{\theta_1}$. We shall call the AF -algebra $\mathbb{A}$ a *toric AF -algebra*, whenever $g \geq 2$.

5. Main result.

Theorem 1 *Let $V = Ob(\mathcal{S})$ and $W = Ob(\mathcal{A}^*)$. The map F^* satisfies the following properties:*

(i) $V \cong W \times (0, \infty)$ is a trivial fiber bundle, whose projection map $\pi : V \rightarrow W$ coincides with F^* ;

(ii) F^* is a covariant functor from the category $\mathcal{S}$ to the category $\mathcal{A}^*$, which maps isomorphic Riemann surfaces $\mathbb{S}, \mathbb{S}' \in \mathcal{S}$ to the stably isomorphic toric AF -algebras $\mathbb{A}, \mathbb{A}' \in \mathcal{A}^*$.

6. Structure of the paper.

The article is organized as follows. In section 2, the toric AF -algebras and functor F^* are (re-)introduced. The main result (theorem 1) is proved in section 3. As a corollary, one gets a functor between the complex tori and Effros-Shen algebras. Finally, section 4 is an appendix, where a brief review of the preliminary result can be found.

2 Notation

In the present section we introduce an object, which we shall call a toric AF -algebra. This is a special type of the AF -algebra, which can be regarded as a generalization of the Effros-Shen algebra to the case $g \geq 2$.

2.1 The toric AF -algebras

The following is a generalization of the Effros-Shen algebras to the case of genus $g \geq 2$. We shall call the object a toric AF -algebra¹.

1. Definition

Let

$$\begin{pmatrix} 1 \\ \theta_1 \\ \vdots \\ \theta_{n-1} \end{pmatrix} = \lim_{k \rightarrow \infty} \prod_{i=1}^k \begin{pmatrix} 0 & 0 & \dots & 0 & 1 \\ 1 & 0 & \dots & 0 & b_1^{(i)} \\ \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & \dots & 1 & b_{n-1}^{(i)} \end{pmatrix} \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix},$$

where

$$n = \begin{cases} 6g - 6, & \text{if } g \geq 2 \\ 2, & \text{if } g = 1, \end{cases}$$

be a *convergent* Jacobi-Perron continued fraction. Let $\mathbb{A} := \mathbb{A}_{\theta_1, \dots, \theta_{n-1}}$ be an AF -algebra given by the Bratteli diagram in Fig.2 (drawn in the case $g = 2$); the $b_j^{(i)}$ are the multiplicity of edges of the graph. The AF -algebra $\mathbb{A}$ will be called a *toric AF -algebra* of genus $g \geq 2$.

2. Equivalent definition

In view of Bauer's criterion (§4.3.4), one can define a toric AF -algebra as an AF -algebra with a unique trace. Namely, the following lemma is true.

¹I am indebted to Yu. I. Manin for the term.




Figure 2: The toric AF -algebra of genus $g = 2$.

Lemma *The AF -algebra $\mathbb{A}$ is a toric AF -algebra if and only if it has a unique trace.*

Proof. Let $\mathbb{A}$ be a toric AF -algebra. Then the convergence of the corresponding Jacobi-Perron continued fraction implies the unique trace, see an argument found in the proof of Theorem 4 of Bauer's paper [1].

The converse, that the unique trace implies convergence of the Jacobi-Perron fraction, is rather direct consequence of the definitions, or one may argue as in the proof of Theorem 3.2 of Effros-Shen [6]. $\square$

3. Examples

We conclude with two examples, which illustrate a difference between the toric AF -algebras and general AF -algebras. One example is based on the Perron sufficient conditions for a convergence of the Jacobi-Perron fractions, another (a counterexample) on the Effros-Shen calculations made in [6].

Example 1. Let $\mathbb{A} = \mathbb{A}_{\theta_1, \dots, \theta_{n-1}}$ be an AF -algebra, whose Jacobi-Perron continued fraction satisfies the conditions $0 < \frac{1}{b_{n-1}^{(k)}} \leq C$ and $0 \leq \frac{b_i^{(k)}}{b_{n-1}^{(k)}} \leq C$, where C is a real constant independent of k . Then, by Perron's sufficient conditions (§4.3.5), the continued fraction is convergent. Thus, the AF -algebra $\mathbb{A}$ is a toric AF -algebra.

Example 2. Let $n = 3$ and the AF -algebra $\mathbb{A} := \mathbb{A}_{\theta_1, \theta_2}$ has the Jacobi-Perron

continued fraction of the form:

$$\lim_{k \rightarrow \infty} \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & \beta_1 \\ 0 & 1 & 0 \end{pmatrix} \dots \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & \beta_k \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix},$$

where $\sum_{k=1}^{\infty} \frac{1}{\beta_k} < 1$. It has been shown by Effros and Shen [6] (Proposition 4.1), that the above Jacobi-Perron continued fraction is divergent. Thus, $\mathbb{A}$ is the AF -algebra, which is *not* a toric AF -algebra.

2.2 The map $F^* : Ob(\mathcal{S}) \rightarrow Ob(\mathcal{A}^*)$

1. Category $\mathcal{S}$.

Let $T(g)$ be the Teichmüller space of genus $g \geq 1$. By the Hubbard-Masur theory, $T(g) - \{pt\} \cong \Phi_X$, where Φ_X is the space of measured foliations on a surface of genus g . Recall that the HM -foliation is a measured foliation, whose regular leaves are dense in X . By $V \subset T(g)$ we shall understand a preimage under homomorphism $h : T(g) - \{pt\} \rightarrow \Phi_X$ of the set of HM -foliations such that for any $\mathcal{F} \in V$:

- (i) *Sing* $\mathcal{F}$ consists of singular points, which are:
 - (a) the singular points with $k = 1$ (tripods), if $g \geq 2$, or
 - (b) the singular points with $k = 0$ (fake saddles), if $g = 1$;
- (ii) $\mathcal{F}$ is a uniquely ergodic measured foliation.

Lemma *The V is a generic subset of $T(g)$.*

Proof. Let $g \geq 2$ and $h(V^i)$ a subset of Φ_X satisfying condition (i). Singular points of degree 1 correspond to simple zeroes of holomorphic quadratic forms on X , endowed with a complex structure. Holomorphic forms with such zeroes are stable with respect to small perturbation in the space of holomorphic quadratic forms and, in view of Morse theory, make a generic set therein. Since h is continuous, V^i is a generic set of $T(g)$. In case $g = 1$, observe that fake saddle is a removable singularity, i.e. the corresponding holomorphic form has no singular points. In this case $h(V^i)$ coincides with the space of all HM -foliations, which are generic on torus for obvious reasons.

Let $h(V^{ii})$ be a subset of Φ_X satisfying condition (ii). It has been shown by Masur [17] and Veech [23] that the measured foliations with a uniquely ergodic invariant measure are generic in Φ_X . Therefore, by continuity of h , V^{ii} is generic in $T(g)$. It has been shown in the above cited works, that $h(V^{ii})$ is generic in $h(V^i)$. Let $V = V^i \cap V^{ii}$. Since a generic subset of generic set is generic in $T(g)$, lemma follows. $\square$

Definition 1 *Let $Ob(\mathcal{S}) = \{\mathbb{S} \in V \mid \varphi(\mathbb{S}) \in V, \varphi \in Mod(X)\}$ be a maximal subset of V closed under the isomorphisms of Riemann surfaces. The category $\mathcal{S} = Ob(\mathcal{S}) \cup Mor(\mathcal{S})$, where $Mor(\mathcal{S})$ are isomorphisms between the Riemann surfaces $\mathbb{S} \in Ob(\mathcal{S})$.*

2. Category $\mathcal{A}^*$.

Denote by $NCS(g)$ a set of the toric AF -algebras of genus g , where $g \geq 1$ and

$$n = \begin{cases} 6g - 6, & \text{if } g \geq 2 \\ 2, & \text{if } g = 1. \end{cases}$$

Definition 2 Let $Ob(\mathcal{A}^*) = \{\mathbb{A}_{\theta_1, \dots, \theta_{n-1}} \in NCS(g) \mid \text{if } \mathbb{A}_{\theta'_1, \dots, \theta'_{n-1}} \cong \mathbb{A}_{\theta_1, \dots, \theta_{n-1}}, \text{ then } \mathbb{A}_{\theta'_1, \dots, \theta'_{n-1}} \in NCS(g)\}$ be maximal subset of $NCS(g)$ closed under the stable isomorphisms between the AF -algebras. The category $\mathcal{A}^* = Ob(\mathcal{A}^*) \cup Mor(\mathcal{A}^*)$, where $Mor(\mathcal{A}^*)$ are stable isomorphisms between the AF -algebras $\mathbb{A}_{\theta_1, \dots, \theta_{n-1}} \in Ob(\mathcal{A}^*)$.

Note that $\mathcal{A}^* = \mathcal{A}$ whenever $g = 1$.

3. Mapping $F^* : Ob(\mathcal{S}) \rightarrow Ob(\mathcal{A}^*)$.

Let $\mathbb{S} \in Ob(\mathcal{S})$ be a Riemann surface of genus $g \geq 1$. The assignment $\mathbb{S} \rightarrow \mathbb{A}_{\theta_1, \dots, \theta_{n-1}}$, where $\mathbb{A}_{\theta_1, \dots, \theta_{n-1}} \in \mathcal{A}^*$, is as follows. Let $h : T(g) - \{pt\} \rightarrow \Phi_X$ be the Hubbard-Masur homeomorphism. Then $h(\mathbb{S}) = \mathcal{F}(\lambda_1, \dots, \lambda_n)$, where $(\lambda_1, \dots, \lambda_n)$ are the Douady-Hubbard coordinates in Φ_X . Without loss of generality, we assume $\lambda_1 \neq 0$ and let $\theta_1 = \frac{\lambda_1}{\lambda_1}, \dots, \theta_{n-1} = \frac{\lambda_n}{\lambda_1}$.

Lemma $\mathbb{A}_{\theta_1, \dots, \theta_{n-1}}$ is a toric AF -algebra.

Proof. Indeed, by item (i) of definition of the category $\mathcal{S}$, all singular points of the foliation $\mathcal{F}$ are the tripods (simple zeroes). In this case, the argument of [4] shows that $\mathcal{F}$ depends on $n = 6g - 6$ ($n = 2$) real parameters if $g \geq 2$ ($g = 1$).

On the other hand, by item (ii) of the definition of the category $\mathcal{S}$, $\mathcal{F}$ is a uniquely ergodic HM -foliation. By Bauer's criterion, the Jacobi-Perron continued fraction for the vector $(\theta_1, \dots, \theta_{n-1})$ is convergent. Thus, $\mathbb{A}_{\theta_1, \dots, \theta_{n-1}}$ is a toric AF -algebra. $\square$

Definition 3 By $F^* : Ob(\mathcal{S}) \rightarrow Ob(\mathcal{A}^*)$, we understand a map given by the formula $\mathbb{S} \mapsto \mathbb{A}_{\theta_1, \dots, \theta_{n-1}}$.

3 Proof of theorem 1

Let us outline the proof. We shall consider the following sets of objects:

- (i) generic Riemann surfaces V ;
- (ii) pseudo-lattices ² $\mathcal{PL}$;
- (iii) projective pseudo-lattices $\mathcal{PPL}$;
- (iv) the toric AF -algebras W .

²The terminology is due to Yu. I. Manin, who introduced pseudo-lattices in the context of the real multiplication [15].

The proof takes the following steps:

- (a) to show that $V \cong \mathcal{PL}$ are equivalent categories such that isomorphic Riemann surfaces $\mathbb{S}, \mathbb{S}' \in V$ map to isomorphic pseudo-lattices $PL, PL' \in \mathcal{PL}$;
- (b) a non-injective functor $F : \mathcal{PL} \rightarrow \mathcal{PPL}$ is constructed. The F maps isomorphic pseudo-lattices to isomorphic projective pseudo-lattices and $\text{Ker } F \cong (0, \infty)$;
- (c) to show that a subcategory $U \subseteq \mathcal{PPL}$ and W are the equivalent categories.

In other words, we have the following diagram:

$$V \xrightarrow{\alpha} \mathcal{PL} \xrightarrow{F} U \xrightarrow{\beta} W, \quad (1)$$

where α is an injective map, β is a bijection and $\text{Ker } F \cong (0, \infty)$.

Category V . A *Riemann surface* is a triple $(X, \mathbb{S}, j)$, where X is a topological surface of genus $g \geq 1$, $j : X \rightarrow \mathbb{S}$ is a complex (conformal) parametrization of X and $\mathbb{S}$ is a Riemann surface. A *morphism* of Riemann surfaces $(X, \mathbb{S}, j) \rightarrow (X, \mathbb{S}', j')$ is a biholomorphic map modulo the ones, which are isotopic to the identity map with respect to a fixed topological marking of X . A category of generic Riemann surfaces $\mathcal{S}$, consists of $Ob(\mathcal{S})$ which are Riemann surfaces $\mathbb{S} \in V \subset T(g)$ and morphisms $H(\mathbb{S}, \mathbb{S}')$ between $\mathbb{S}, \mathbb{S}' \in Ob(V)$ which coincide with the morphisms specified above. For any $\mathbb{S}, \mathbb{S}', \mathbb{S}'' \in Ob(\mathcal{S})$ and any morphisms $\varphi' : \mathbb{S} \rightarrow \mathbb{S}'$, $\varphi'' : \mathbb{S}' \rightarrow \mathbb{S}''$ a morphism $\phi : \mathbb{S} \rightarrow \mathbb{S}''$ is the composite of φ' and φ'' , which we write as $\phi = \varphi''\varphi'$. The identity morphism, $1_{\mathbb{S}}$, is a morphism $H(\mathbb{S}, \mathbb{S})$.

Category $\mathcal{PL}$. A *pseudo-lattice* (of rank n) is a triple $(\Lambda, \mathbb{R}, j)$, where $\Lambda \cong \mathbb{Z}^n$ and $j : \Lambda \rightarrow \mathbb{R}$ is a homomorphism. A morphism of pseudo-lattices $(\Lambda, \mathbb{R}, j) \rightarrow (\Lambda, \mathbb{R}, j')$ is a commutative diagram:

$$\begin{array}{ccc} \mathbb{Z}^n & \xrightarrow{j} & \mathbb{R} \\ \downarrow \varphi & & \downarrow \psi \\ \mathbb{Z}^n & \xrightarrow{j'} & \mathbb{R} \end{array}$$

where φ is a group homomorphism and ψ is an inclusion map, i.e. $j'(\Lambda') \subseteq j(\Lambda)$. Any isomorphism class of a pseudo-lattice contains a representative given by $j : \mathbb{Z}^n \rightarrow \mathbb{R}$ such that

$$j(1, 0, \dots, 0) = \lambda_1, \quad j(0, 1, \dots, 0) = \lambda_2, \quad \dots, \quad j(0, 0, \dots, 1) = \lambda_n,$$

where $\lambda_1, \lambda_2, \dots, \lambda_n$ are positive reals. The pseudo-lattices of rank n make up a category, which we denote by $\mathcal{PL}_n$.

The following lemma implies a functoriality of the map F^* . Roughly, the lemma says that the $\mathbb{Z}$ -module $\mathbb{Z}\lambda_1 + \dots + \mathbb{Z}\lambda_n$ is an invariant of the isomorphism class of the Riemann surface $\mathbb{S}$ (a modulus of $\mathbb{S}$). The action of $Mod(X)$ on such a module corresponds to a transformation of the basis of the module. The proof of lemma is based on the explicit formulas for λ_i .

Lemma 1 (Basic lemma) *Let $g \geq 2$ ($g = 1$) and $n = 6g - 6$ ($n = 2$). There exists an injective covariant functor $\alpha : V \rightarrow \mathcal{PL}_n$ which maps isomorphic Riemann surfaces $\mathbb{S}, \mathbb{S}' \in V$ to the isomorphic pseudo-lattices $PL, PL' \in \mathcal{PL}_n$.*

Proof. Let $\alpha : T(g) - \{pt\} \rightarrow Hom(H_1^{odd}(\tilde{\mathbb{S}}); \mathbb{R}) - 0$ be a Hubbard-Masur map. Since α is a homeomorphism between the respective spaces, we conclude that α is an injective map. The first claim of lemma is proved.

Let us show that α sends morphisms of $\mathcal{S}$ to morphisms of $\mathcal{PL}$. Let $\varphi \in Mod(X)$ be a diffeomorphism of X . Suppose that all the zeroes of measured foliations are generic (simple) and let $p : \tilde{X} \rightarrow X$ be the double cover of X . (Note that the case of torus does not require a double cover, and thus one can assert $p = Id$ in the argument below.) Denote by $\tilde{\varphi}$ a diffeomorphism of $\tilde{X}$, which makes the following diagram commutative:

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\tilde{\varphi}} & \tilde{X} \\ \downarrow p & & \downarrow p \\ X & \xrightarrow{\varphi} & X \end{array}$$

One can consider the effect of $\varphi, \tilde{\varphi}$ and p on the respective (relative) integral homology groups:

$$\begin{array}{ccc} H_1^{odd}(\tilde{X}) \oplus H_1^{even}(\tilde{X}) & \xrightarrow{\tilde{\varphi}_*} & H_1^{odd}(\tilde{X}) \oplus H_1^{even}(\tilde{X}) \\ \downarrow p_* & & \downarrow p_* \\ H_1(X, Sing \mathcal{F}) & \xrightarrow{\varphi_*} & H_1(X, Sing \mathcal{F}) \end{array}$$

where $Ker p_* \cong H_1^{even}(\tilde{X})$. Since $p_* : H_1^{odd}(\tilde{X}) \rightarrow H_1(X, Sing \mathcal{F})$ is an isomorphism, we conclude that $\tilde{\varphi}_* \in GL_n(\mathbb{Z})$, where $n = \dim H_1^{odd}(\tilde{X})$. It is easy to see, that $\tilde{\varphi}_*$ acts on a pseudo-lattice by a transformation of its basis, and therefore, $\tilde{\varphi}_* \in Mor(\mathcal{PL})$.

Let us show that α is a functor. Indeed, let $\mathbb{S}, \mathbb{S}' \in V$ be isomorphic Riemann surfaces, such that $\mathbb{S}' = \varphi(\mathbb{S})$ for a $\varphi \in \text{Mod}(X)$. Let a_{ij} be the elements of matrix $\tilde{\varphi}_* \in GL_n(\mathbb{Z})$. Recall that:

$$\lambda_i = \int_{\gamma_i} \phi \quad (2)$$

for a closed 1-form $\phi = \text{Re } \omega$ and $\gamma_i \in H_1^{\text{odd}}(\tilde{X})$. Then

$$\gamma_j = \sum_{i=1}^n a_{ij} \gamma_i, \quad j = 1, \dots, n, \quad (3)$$

are the elements of a new basis in $H_1^{\text{odd}}(\tilde{X})$. By the integration rules,

$$\lambda'_j = \int_{\gamma_j} \phi = \int_{\sum a_{ij} \gamma_i} \phi = \sum_{i=1}^n a_{ij} \lambda_i. \quad (4)$$

Finally, let $j(\Lambda) = \mathbb{Z}\lambda_1 + \dots + \mathbb{Z}\lambda_n$ and $j'(\Lambda) = \mathbb{Z}\lambda'_1 + \dots + \mathbb{Z}\lambda'_n$. Since $\lambda'_j = \sum_{i=1}^n a_{ij} \lambda_i$ and $(a_{ij}) \in GL_n(\mathbb{Z})$, we conclude that:

$$j(\Lambda) = j'(\Lambda). \quad (5)$$

In other words, the $\mathbb{Z}$ -module $\mathbb{Z}\lambda_1 + \dots + \mathbb{Z}\lambda_n$ is an invariant of $\text{Mod}(X)$. In particular, the pseudo-lattices $(\Lambda, \mathbb{R}, j)$ and $(\Lambda, \mathbb{R}, j')$ are isomorphic. Hence, $\alpha : V \rightarrow \mathcal{P}\mathcal{L}$ maps isomorphic Riemann surfaces to the isomorphic pseudo-lattices, i.e. α is a functor.

Finally, let us show that α is a covariant functor. Indeed, let $\varphi_1, \varphi_2 \in \text{Mor}(\mathcal{S})$. Then $\alpha(\varphi_1 \varphi_2) = (\widetilde{\varphi_1 \varphi_2})_* = (\tilde{\varphi}_1)_* (\tilde{\varphi}_2)_* = \alpha(\varphi_1) \alpha(\varphi_2)$. Lemma 1 follows. $\square$

Category $\mathcal{P}\mathcal{P}\mathcal{L}$. A *projective pseudo-lattice* (of rank n) is a triple $(\Lambda, \mathbb{R}, j)$, where $\Lambda \cong \mathbb{Z}^n$ and $j : \Lambda \rightarrow \mathbb{R}$ is a homomorphism. A morphism of projective pseudo-lattices $(\Lambda, \mathbb{C}, j) \rightarrow (\Lambda, \mathbb{R}, j')$ is a commutative diagram:

$$\begin{array}{ccc} \mathbb{Z}^n & \xrightarrow{j} & \mathbb{R} \\ \downarrow \varphi & & \downarrow \psi \\ \mathbb{Z}^n & \xrightarrow{j'} & \mathbb{R} \end{array}$$

where φ is a group homomorphism and ψ is an $\mathbb{R}$ -linear map. (Notice that unlike the case of pseudo-lattices, ψ is a scaling map as opposite to an inclusion map. This allows to the two pseudo-lattices to be projectively equivalent, while

being distinct in the category $\mathcal{PL}_n$.) It is not hard to see that any isomorphism class of a projective pseudo-lattice contains a representative given by $j : \mathbb{Z}^n \rightarrow \mathbb{R}$ such that

$$j(1, 0, \dots, 0) = 1, \quad j(0, 1, \dots, 0) = \theta_1, \quad \dots, \quad j(0, 0, \dots, 1) = \theta_{n-1},$$

where θ_i are positive reals. The projective pseudo-lattices of rank n make up a category, which we denote by $\mathcal{PPL}_n$.

Category W . Finally, the toric AF -algebras $\mathbb{A} = \mathbb{A}_{\theta_1, \dots, \theta_{n-1}}$, modulo the stable isomorphism between them, make up a category, which we shall denote by W_n .

The following equivalence of categories is immediate.

Lemma 2 *Let $U_n \subseteq \mathcal{PPL}_n$ be a subcategory consisting of the projective pseudo-lattices $PPL = PPL(1, \theta_1, \dots, \theta_{n-1})$ for which the Jacobi-Perron fraction of the vector $(1, \theta_1, \dots, \theta_{n-1})$ converges to the vector. Define a map $\beta : U_n \rightarrow W_n$ by the formula*

$$PPL(1, \theta_1, \dots, \theta_{n-1}) \mapsto \mathbb{A}_{\theta_1, \dots, \theta_{n-1}}.$$

Then β is a bijective functor, which maps isomorphic projective pseudo-lattices to the stably isomorphic toric AF -algebras.

Proof. It is evident that β is injective and surjective. Let us show that β is a functor. Indeed, according to §4.2.4, every totally ordered abelian group of rank n has form $\mathbb{Z} + \theta_1 \mathbb{Z} + \dots + \mathbb{Z} \theta_{n-1}$. The latter is a projective pseudo-lattice PPL from the category U_n . On the other hand, by the Elliott theorem (§4.2.5.C), the PPL defines a stable isomorphism class of the AF -algebra $\mathbb{A}_{\theta_1, \dots, \theta_{n-1}} \in W_n$. Therefore, β maps isomorphic projective pseudo-lattices (from the set U_n) to the stably isomorphic toric AF -algebras, and *vice versa*. Lemma 2 follows. $\square$

Let $PL(\lambda_1, \lambda_2, \dots, \lambda_n) \in \mathcal{PL}_n$ and $PPL(1, \theta_1, \dots, \theta_{n-1}) \in \mathcal{PPL}_n$. To finish the proof of theorem 1, it remains to show the following.

Lemma 3 *Let $F : \mathcal{PL}_n \rightarrow \mathcal{PPL}_n$ be a map given by formula*

$$PL(\lambda_1, \lambda_2, \dots, \lambda_n) \mapsto PPL\left(1, \frac{\lambda_2}{\lambda_1}, \dots, \frac{\lambda_n}{\lambda_1}\right).$$

Then $\text{Ker } F = (0, \infty)$ and F is a functor which maps isomorphic pseudo-lattices to isomorphic projective pseudo-lattices.

Proof. Indeed, F can be thought as a map from $\mathbb{R}^n$ to $\mathbb{R}P^{n-1}$. Hence $\text{Ker } F = \{\lambda_1 : \lambda_1 > 0\} \cong (0, \infty)$. The second part of lemma is evident. $\square$

Assuming $n = 6g - 6$ ($n = 2$) for $g \geq 2$ ($g = 1$), one gets theorem 1 from lemmas 1-3. $\square$

Recall that our study has been motivated by a correspondence between the complex tori and the Effros-Shen algebras. In view of theorem 1, we have the following corollary.

Corollary 1 *There exists a covariant (non-injective) functor $F : \mathcal{B} \rightarrow \mathcal{A}$, which maps isomorphic complex tori to the stably isomorphic Effros-Shen algebras.*

Proof. Let $g = 1$ and $n = 2$ in §3, and repeat the argument. $\square$

4 Appendix

The present section contains a minimal background information used to prove our main results. The measured foliations are expounded in [9], [11] and [22]. A full exposition of the remarkable connection between measured foliations and the Teichmüller spaces (Hubbard-Masur theory) can be found in [11]. For the C^* -algebra part, especially AF -algebras, we refer the reader to [5]. Finally, a comprehensive account of the Jacobi-Perron continued fractions can be found in [2].

4.1 Hubbard-Masur theory

1. Hubbard-Masur homeomorphism

Let $T(g)$ be the Teichmüller space of the topological surface X of genus $g \geq 1$, i.e. the space of the complex (conformal) structures on X . Consider the vector bundle $p : Q \rightarrow T(g)$ over $T(g)$, whose fiber above the point $\mathbb{S} \in T(g)$ is a vector space $H^0(\mathbb{S}, \Omega^{\otimes 2})$. Given a non-zero $q \in Q$ above $\mathbb{S}$, we can consider a horizontal measured foliation $\mathcal{F}_q \in \Phi_X$, where Φ_X denotes the space of the equivalence classes of measured foliations on X . If $\{0\}$ is the zero section of Q , the above construction defines a map $Q - \{0\} \rightarrow \Phi_X$. For any $\mathcal{F} \in \Phi_X$, let $E_{\mathcal{F}} \subset Q - \{0\}$ be the fiber above $\mathcal{F}$. In other words, $E_{\mathcal{F}}$ is a subspace of the holomorphic quadratic forms, whose horizontal trajectory structure coincides with the measured foliation $\mathcal{F}$.

Theorem ([11]) *The restriction $E_{\mathcal{F}} \rightarrow T(g)$ of p to $E_{\mathcal{F}}$ is a homeomorphism.*

2. Parametrization of the Teichmüller space by measured foliations

One obtains an important corollary from the Hubbard-Masur theorem. We shall assume that $\mathcal{F} \in \Phi_X$ is a measured foliation, whose set of singular points consists of simple zeroes (i.e. $k_1 = 1$ for all the singular points). Denote by $\mathcal{F}'$ a measured foliation defined by the vertical trajectories $\operatorname{Re} q = 0$, where $q \in E_{\mathcal{F}}$. Since $\mathcal{F} = \operatorname{Const}$ for all $q \in E_{\mathcal{F}}$, we conclude that $\mathcal{F}' \in \Phi_X$ parametrizes the space $E_{\mathcal{F}}$. The $\mathcal{F}'$ and $\mathcal{F}$ have a common set of the singular points, and therefore all zeroes of $\mathcal{F}'$ are simple. Thus, $\mathcal{F}'$ varies in a subset of Φ_X of dimension $n = 6g - 6$ ($n = 2$) if $g \geq 2$ ($g = 1$). Recall that $\Phi_X \cong \mathbb{R}^n - 0$, while $T(g) \cong \mathbb{R}^n$. We let pt be the image of the zero quadratic differential under the Hubbard-Masur homeomorphism.

Corollary *The mapping $h_{E_{\mathcal{F}}} : \Phi_X \rightarrow T(g) - \{pt\}$ is a homeomorphism.*

3. *The canonical homeomorphism $h : \Phi_X \rightarrow T(g) - \{pt\}$.*

It follows from the Hubbard-Masur construction, that the homeomorphism $h_{E_{\mathcal{F}}}$ depends on an initial choice of the measured foliation $\mathcal{F}$. Let us show, however, that there exists a canonical map $h : \Phi_X \rightarrow T(g) - \{pt\}$. Let $\mathcal{S}$ be the set of homotopy equivalence classes of simple closed curves on the surface X . Define a map $i : \Phi_X \rightarrow \mathbb{R}^{\mathcal{S}}$ by the formula

$$\phi \mapsto \inf_{\gamma_i} \int_{\gamma_i} |\phi(t)| \gamma'(t) dt, \quad \phi \in \Phi_X, \quad \gamma_i \in \mathcal{S},$$

where the infimum is taken over all simple closed curves in the homotopy class γ_i . Recall that a *length spectrum* of a Riemann surface $\mathbb{S} \in T(g)$ is an element of $\mathbb{R}^{\mathcal{S}}$ consisting of the lengths of the shortest simple closed geodesic representatives in each of the homotopy classes of $\mathcal{S}$. We shall denote the latter function by $j : T(g) \rightarrow \mathbb{R}^{\mathcal{S}}$. The homeomorphism $h : \Phi_X \rightarrow T(g) - \{pt\}$ will be called *canonical* if the following diagram is commutative:

$$\begin{array}{ccc} \Phi_X & \xrightarrow{h} & T(g) \\ & \searrow i & \downarrow j \\ & & \mathbb{R}^{\mathcal{S}} \end{array}$$

Further, by a *Hubbard-Masur homeomorphism* we shall understand the canonical homeomorphism h .

4. *Douady-Hubbard coordinates in the space of measured foliations*

A. Recall that Φ_X is the space of equivalence classes of measured foliations on the topological surface X . Following Douady and Hubbard [4], we consider a coordinate system on Φ_X , suitable for the proof of theorem 1. For clarity, let us make a generic assumption that $q \in H^0(\mathbb{S}, \Omega^{\otimes 2})$ is a non-trivial holomorphic quadratic differential with only simple zeroes. We wish to construct a Riemann surface of $\sqrt{q}$, which is a double cover of $\mathbb{S}$ with ramification over the zeroes of q . Such a surface, denoted by $\tilde{\mathbb{S}}$, is unique and has an advantage of carrying a holomorphic differential ω , such that $\omega^2 = q$. We further denote by $\pi : \tilde{\mathbb{S}} \rightarrow \mathbb{S}$ the covering projection. The vector space $H^0(\tilde{\mathbb{S}}, \Omega)$ splits into the direct sum $H_{even}^0(\tilde{\mathbb{S}}, \Omega) \oplus H_{odd}^0(\tilde{\mathbb{S}}, \Omega)$ in view of the involution π^{-1} of $\tilde{\mathbb{S}}$, and the vector space:

$$H^0(\mathbb{S}, \Omega^{\otimes 2}) \cong H_{odd}^0(\tilde{\mathbb{S}}, \Omega).$$

Let $H_1^{odd}(\tilde{\mathbb{S}})$ be the odd part of the homology of $\tilde{\mathbb{S}}$ relatively the zeroes of q . Consider the pairing $H_1^{odd}(\tilde{\mathbb{S}}) \times H^0(\mathbb{S}, \Omega^{\otimes 2}) \rightarrow \mathbb{C}$, defined by the integration $(\gamma, q) \mapsto \int_{\gamma} \omega$. We shall take the associated map $\psi_q : H^0(\mathbb{S}, \Omega^{\otimes 2}) \rightarrow \text{Hom}(H_1^{odd}(\tilde{\mathbb{S}}); \mathbb{C})$ and let $h_q = \text{Re } \psi_q$.

Lemma ([4]) *The map*

$$h_q : H^0(\mathbb{S}, \Omega^{\otimes 2}) \longrightarrow \text{Hom} (H_1^{\text{odd}}(\tilde{\mathbb{S}}); \mathbb{R}),$$

is an $\mathbb{R}$ -isomorphism.

B. Since each $\mathcal{F} \in \Phi_X$ is the vertical foliation $Re\, q = 0$ for a $q \in H^0(\mathbb{S}, \Omega^{\otimes 2})$, the Douady-Hubbard lemma implies that $\Phi_X \cong \text{Hom} (H_1^{\text{odd}}(\tilde{\mathbb{S}}); \mathbb{R})$. By formulas for the relative homology, one finds that $H_1^{\text{odd}}(\tilde{\mathbb{S}}) \cong \mathbb{Z}^n$, where

$$n = \begin{cases} 6g - 6, & \text{if } g \geq 2 \\ 2, & \text{if } g = 1. \end{cases}$$

Finally, each $h \in \text{Hom} (\mathbb{Z}^n; \mathbb{R})$ is given by the reals $\lambda_1 = h(e_1), \dots, \lambda_n = h(e_n)$, where $(e_1, \dots, e_n)$ is a basis in $\mathbb{Z}^n$. Thus, the reals $(\lambda_1, \dots, \lambda_n)$ make a coordinate system in the linear space Φ_X .

4.2 AF-algebras

1. C^* -algebras

A. Rings of bounded operators on Hilbert space have been introduced in the 1930s by Murray and von Neumann. These rings, called von Neumann algebras, have a plethora of remarkable properties, with deep and unexpected ties to geometry, topology and representation theory.

B. By the C^* -algebra one understands a noncommutative Banach algebra with an involution. Namely, a C^* -algebra A is an algebra over $\mathbb{C}$ with a norm $a \mapsto \|a\|$ and an involution $a \mapsto a^*$, $a \in A$, such that A is complete with respect to the norm, and such that $\|ab\| \leq \|a\| \|b\|$ and $\|a^*a\| = \|a\|^2$ for every $a, b \in A$. Every C^* -algebra is isomorphic to a subalgebra of algebra of bounded linear operators on a Hilbert space.

C. If A is commutative, then the Gelfand theorem says that A is isomorphic to the C^* -algebra $C_0(X)$ of continuous complex-valued functions on a locally compact Hausdorff space X . For otherwise, A represents a noncommutative topological space X .

2. Dimension groups

Let A be a unital C^* -algebra and $V(A)$ be the union (over n) of projections in the $n \times n$ matrix C^* -algebra with entries in A . Projections $p, q \in V(A)$ are equivalent if there exists a partial isometry u such that $p = u^*u$ and $q = uu^*$. The equivalence class of projection p is denoted by $[p]$. The equivalence classes of orthogonal projections can be made to a semigroup by putting $[p] + [q] = [p \oplus q]$. The Grothendieck completion of this semigroup to an abelian group is called a K_0 -group of algebra A . The functor $A \rightarrow K_0(A)$ maps a category of unital C^* -algebras into the category of abelian groups so that projections in algebra

A correspond to a positive cone $K_0^+ \subset K_0(A)$ and the unit element $1 \in A$ corresponds to an order unit $u \in K_0(A)$. The ordered abelian group (K_0, K_0^+, u) with an order unit is called a *dimension group*.

3. Isomorphism of dimension groups

We will need to define an isomorphism for the dimension groups. Recall that by an ordered group one understands an abelian group G together with a subset $P = G^+$ such that $P + P \subseteq P$, $P \cap (-P) = \{0\}$, and $P - P = G$. We call P the positive cone on G . We write $a \leq b$ (or $a < b$) if $b - a \in P$ (or $b - a \in P \setminus \{0\}$). Given ordered groups G and H , we say that a homomorphism $\varphi : G \rightarrow H$ is positive if $\varphi(G^+) \subseteq H^+$, and that $\varphi : G \rightarrow H$ an order-isomorphism if $\varphi(G^+) = H^+$. An *isomorphism* $\cong$ between dimension groups $(K_0(A), K_0^+(A), u)$ and $(K_0(A'), K_0^+(A'), u')$ is an order-isomorphism such that $\varphi(u) = u'$.

4. Dimension groups as dense subgroups of $\mathbb{R}$

A positive homomorphism $s : G \rightarrow \mathbb{R}$ is called a *state* if $s(u) = 1$, where $u \in G^+$ is an order unit of G . We let $S(G)$ be the state space of G , i.e. the set of states on G endowed with natural topology. $S(G)$ is a compact convex subset of vector space $\text{Hom}(G; \mathbb{R})$. By the Krein-Milman theorem, $S(G)$ is the closed convex hull of its extreme points. When $S(G)$ consists of one point, the abelian group G is said to be *totally ordered*. The totally ordered abelian groups of rank $n \geq 2$ will be of main interest for us, and the standard way to produce such groups is this. Let $G \subset \mathbb{R}$ be a dense abelian subgroup of rank n . The positive cone of G is defined to be the set $G^+ = \{g \in G : g > 0\}$ and the order unit $u = 1$. Any totally ordered abelian group appears in this way [5], Corollary 4.7.

5. The AF -algebras, Bratteli diagrams and Elliott invariants

A. One of the major classes of the C^* -algebras whose structure and invariants are well-understood, are the AF -algebras. Recall that every finite-dimensional C^* -algebra is isomorphic to $A = M_{n_1}(\mathbb{C}) \oplus \dots \oplus M_{n_r}(\mathbb{C})$, where $n_1, \dots, n_r$ are positive integers and $M_{n_k}(\mathbb{C})$ are the C^* -algebras of square matrices with the complex entries. If A, B are finite-dimensional C^* -algebras, then $\text{Hom}(A, B)$ are described by systems of the matrix units in A, B . Namely, the matrix unit $E_{ij}^{n_k}$ in A is a matrix whose entries are zero except $e_{ij} = 1$. For very obvious reasons, the matrix units of A is a linear basis of A over $\mathbb{C}$. If $h \in \text{Hom}(A, B)$, then h is defined by the image, $h(E_{ij}^{n_k})$, in the set of matrix units of B . If there are other matrix units of A which map to $h(E_{ij}^{n_k})$, we denote by μ their total number. It easy to see that numbers μ are all one needs to describe h . Indeed, since A and B are direct sums of their matrix units, one can rearrange the summands in A and B , so that h acts by the formula $\oplus_{j=1}^r \mu_{ij} a_i \mapsto b_j$, see [5] for details of the construction. The AF -algebra, $\mathbb{A}$, is the inductive limit of finite-dimensional C^* -algebras

$$A_1 \xrightarrow{\mu_1} A_2 \xrightarrow{\mu_2} \dots$$

induced by the homomorphisms $\mu_k : A_k \rightarrow A_{k+1}$. Here $\mu_k = (\mu_{ij}^k)$ are the multiplicity matrices specified above.

B. The ∞ -partite graph whose incidence matrix on the k -th step is μ_k , is called a *Bratteli diagram* of $\mathbb{A}$. For an example of such a diagram, see Fig.1. Note that $\mathbb{A}$ is no longer a finite-dimensional C^* -algebra, hence the name *AF* (the approximately finite-dimensional C^* -algebra).

C. Denote by $\mathbb{Z}$ and $\mathbb{Z}_+$ the integers and non-negative integers, respectively. We shall restrict to the *AF*-algebras, such that $\mu_i \in GL_n(\mathbb{Z})$. Since $K_0(A_i) \cong \mathbb{Z}^n$, we conclude that $K_0(\mathbb{A}) \cong \mathbb{Z}^n$. Similarly, $K_0^+(\mathbb{A}) = \lim_{i \rightarrow \infty} \mu_1 \dots \mu_i(\mathbb{Z}_+^n)$. Thus, at least formally, we know the dimension group of $\mathbb{A}$. The following beautiful result describes the isomorphism and stable isomorphism classes of the *AF*-algebras.

Lemma ([8]) *Let $\mathbb{A}, \mathbb{A}'$ be the AF-algebras. Then:*

- (i) $\mathbb{A} \cong \mathbb{A}'$ if and only if $(K_0(\mathbb{A}), K_0^+(\mathbb{A}), u) \cong (K_0(\mathbb{A}'), K_0^+(\mathbb{A}'), u')$;
- (ii) $\mathbb{A} \otimes \mathcal{K} \cong \mathbb{A}' \otimes \mathcal{K}$ if and only if $(K_0(\mathbb{A}), K_0^+(\mathbb{A})) \cong (K_0(\mathbb{A}'), K_0^+(\mathbb{A}'))$.

4.3 The Jacobi-Perron continued fractions

1. Regular continued fractions

Let $a_1, a_2 \in \mathbb{N}$ such that $a_2 \leq a_1$. Recall that the greatest common divisor of a_1, a_2 , $GCD(a_1, a_2)$, can be determined from the Euclidean algorithm:

$$\begin{cases} a_1 &= a_2 b_1 + r_3 \\ a_2 &= r_3 b_2 + r_4 \\ r_3 &= r_4 b_3 + r_5 \\ \vdots & \\ r_{k-3} &= r_{k-2} b_{k-1} + r_{k-1} \\ r_{k-2} &= r_{k-1} b_k, \end{cases}$$

where $b_i \in \mathbb{N}$ and $GCD(a_1, a_2) = r_{k-1}$. The Euclidean algorithm can be written as the regular continued fraction

$$\theta = \frac{a_1}{a_2} = b_1 + \frac{1}{b_2 + \frac{1}{b_3 + \dots + \frac{1}{b_k}}} = [b_1, \dots, b_k].$$

If a_1, a_2 are non-commensurable (i.e. $\theta \in \mathbb{R} - \mathbb{Q}$), then the Euclidean algorithm never stops and $\theta = [b_1, b_2, \dots]$. Note that the regular continued fraction can be written in a (projective) matrix form:

$$\begin{pmatrix} 1 \\ \theta \end{pmatrix} = \lim_{k \rightarrow \infty} \begin{pmatrix} 0 & 1 \\ 1 & b_1 \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ 1 & b_k \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

2. The Jacobi-Perron continued fractions

The Jacobi-Perron algorithm and connected (multidimensional) continued fraction generalizes the Euclidean algorithm to the case $GCD(a_1, \dots, a_n)$ when $n \geq 2$. Namely, let $\lambda = (\lambda_1, \dots, \lambda_n)$, $\lambda_i \in \mathbb{R} - \mathbb{Q}$ and $\theta_{i-1} = \frac{\lambda_i}{\lambda_1}$, where $1 \leq i \leq n$ and $\lambda_1 \neq 0$. Consider a projective equivalence class $(1, \theta_1, \dots, \theta_{n-1})$ of the vector λ .

Definition *The continued fraction*

$$\begin{pmatrix} 1 \\ \theta_1 \\ \vdots \\ \theta_{n-1} \end{pmatrix} = \lim_{k \rightarrow \infty} \begin{pmatrix} 0 & 0 & \dots & 0 & 1 \\ 1 & 0 & \dots & 0 & b_1^{(1)} \\ \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & \dots & 1 & b_{n-1}^{(1)} \end{pmatrix} \cdots \begin{pmatrix} 0 & 0 & \dots & 0 & 1 \\ 1 & 0 & \dots & 0 & b_1^{(k)} \\ \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & \dots & 1 & b_{n-1}^{(k)} \end{pmatrix} \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix},$$

where $b_i^{(j)} \in \mathbb{N} \cup \{0\}$, is called the Jacobi-Perron fraction of the vector $\lambda \in \mathbb{R}^n$.

To recover the integers $b_i^{(k)}$ from the vector $(\theta_1, \dots, \theta_{n-1})$, one has to repeatedly solve the following system of equations:

$$\begin{cases} \theta_1 &= b_1^{(1)} & + \frac{1}{\theta'_{n-1}} \\ \theta_2 &= b_2^{(1)} & + \frac{\theta'_1}{\theta'_{n-1}} \\ \vdots & & \\ \theta_{n-1} &= b_{n-1}^{(1)} & + \frac{\theta'_{n-2}}{\theta'_{n-1}}, \end{cases}$$

where $(\theta'_1, \dots, \theta'_{n-1})$ is the next input vector. Thus, each vector $(\theta_1, \dots, \theta_{n-1})$ gives rise to a formal Jacobi-Perron continued fraction. Whether the fraction is convergent or not, is yet to be determined.

3. Convergent Jacobi-Perron continued fractions

Let us introduce some notation, see p. 13 of [2]. Define $A_i^{(j)} = \delta_{ij}$ to be the Kronecker delta, where $i, j = 0, \dots, n-1$. Suppose that the integers $A_i^{(\nu)}, \dots, A_i^{(\nu+n-1)}$ are known. One can define by the induction:

$$A_i^{(\nu+n)} = \sum_{j=0}^{n-1} b_i^{(\nu)} A_i^{(\nu+j)},$$

where $b_0^{(\nu)} = 1$ for $i = 0, \dots, n-1$ and $\nu = 0, 1, \dots, \infty$.

Definition *The Jacobi-Perron continued fraction is said to be convergent, if $\theta_i = \lim_{k \rightarrow \infty} \frac{A_i^{(k)}}{A_0^{(k)}}$ for all $i = 1, \dots, n-1$.*

Unless $n = 2$, the convergence of the Jacobi-Perron fractions is a delicate question. To the best of our knowledge, there exists no intrinsic necessary and

sufficient conditions for such a convergence. However, the Bauer criterion and Masur-Veech theorem (see below) imply that the Jacobi-Perron fractions converge for a generic set of vectors $\lambda \in \mathbb{R}^n$. We shall give in §4.3.5 an intrinsic sufficient condition, which we shall use to construct the toric AF -algebras.

4. Bauer's criterion

Convergence of the Jacobi-Perron continued fractions can be characterized in terms of measured foliations. Let $\mathcal{F} \in \Phi_X$ be measured foliation on a surface X of genus $g \geq 1$. Recall that $\mathcal{F}$ is called uniquely ergodic if every invariant measure of $\mathcal{F}$ is a multiple of the Lebesgue measure. By the Masur-Veech theorem, there exists a generic subset $V \subset \Phi_X$ such that each $\mathcal{F} \in V$ is uniquely ergodic [17], [23]. We let $\lambda = (\lambda_1, \dots, \lambda_n)$ be the vector with coordinates $\lambda_i = \int_{\gamma_i} \phi$, where $\gamma_i \in H_1^{odd}(\tilde{X})$ and ϕ is the covering 1-form (a flow) of the foliation $\mathcal{F}$ on the covering surface $\tilde{X}$. By an abuse of notation, we shall say that $\lambda \in V$. Note that the covering flow ϕ can be obtained from λ by the method of zippered rectangles; in this case an interval exchange map (λ, π) is defined. The last observation allows us to obtain the following characterization of convergence of the Jacobi-Perron fractions.

Lemma ([1]) *The Jacobi-Perron continued fraction of λ converges if and only if $\lambda \in V \subset \mathbb{R}^n$.*

5. Sufficient conditions of convergence

The following sufficient condition is due to O. Perron.

Lemma ([18], Satz II) *The Jacobi-Perron continued fraction*

$$\lim_{k \rightarrow \infty} \begin{pmatrix} 0 & 0 & \dots & 0 & 1 \\ 1 & 0 & \dots & 0 & b_1^{(1)} \\ \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & \dots & 1 & b_{n-1}^{(1)} \end{pmatrix} \cdots \begin{pmatrix} 0 & 0 & \dots & 0 & 1 \\ 1 & 0 & \dots & 0 & b_1^{(k)} \\ \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & \dots & 1 & b_{n-1}^{(k)} \end{pmatrix} \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix}$$

is convergent, if for all $i = 1, \dots, n-1$ and $k = 1, \dots, \infty$, it holds:

$$0 < \frac{1}{b_{n-1}^{(k)}} \leq C \quad \text{and} \quad 0 \leq \frac{b_i^{(k)}}{b_{n-1}^{(k)}} < C,$$

where C is a real constant independent of k .

Perron's lemma indicates when the continued fractions diverge. In particular, the divergence is caused by a fast growth of the entry $b_i^{(k)}$ with respect to $b_{n-1}^{(k)}$ as $k \rightarrow \infty$. It is easy to see, that the regular continued (i.e. $n = 2$) and periodic Jacobi-Perron fractions are always convergent.

References

- [1] M. Bauer, A characterization of uniquely ergodic interval exchange maps in terms of the Jacobi-Perron algorithm, *Bol. Soc. Bras. Mat.* 27 (1996), 109-128.
- [2] L. Bernstein, The Jacobi-Perron Algorithm, its Theory and Applications, *Lect. Notes in Math.* 207, Springer 1971.
- [3] A. Connes, M. Douglas and A. Schwarz, Noncommutative geometry and matrix theory: compactification on tori, *J. High Energy Phys.* 1998, no. 2, Paper 3, 35 pp. (electronic)
- [4] A. Douady and J. Hubbard, On the density of Strebel differentials, *Inventiones Math.* 30 (1975), 175-179.
- [5] E. G. Effros, Dimensions and C^* -Algebras, in: *Conf. Board of the Math. Sciences, Regional conference series in Math.*, No.46, AMS, 1981.
- [6] E. G. Effros and C.-L. Shen, Dimension groups and finite difference equations, *J. Operator Theory* 2 (1979), 215-231.
- [7] E. G. Effros and C. L. Shen, Approximately finite C^* -algebras and continued fractions, *Indiana Univ. Math. J.* 29 (1980), 191-204.
- [8] G. A. Elliott, On the classification of inductive limits of sequences of semisimple finite-dimensional algebras, *J. Algebra* 38 (1976), 29-44.
- [9] A. Fathi, F. Laudenbach and V. Poénaru, *Travaux de Thurston sur les Surfaces*, Astérisque 66-67, 1979.
- [10] R. Hain and E. Looijenga, Mapping class groups and moduli spaces of curves, *Proc. Symposia in Pure Math.*, vol. 62 (2), 1997, 97-142.
- [11] J. Hubbard and H. Masur, Quadratic differentials and foliations, *Acta Math.* 142 (1979), 221-274.
- [12] F. Klein, *Ausgewählte Kapitel der Zahlentheorie*, *Math. Annalen* 48 (1896/97).
- [13] M. Kontsevich, XI Solomon Lefschetz Memorial Lecture series: Hodge structures in non-commutative geometry. Notes by Ernesto Lupercio. *Contemp. Math.*, 462, Noncommutative geometry in mathematics and physics, 1-21. Amer. Math. Soc., Providence, RI, 2008.
- [14] M. Kontsevich and Y. Soibelman, Notes on A-infinity algebras, A-infinity categories and non-commutative geometry. I, [arXiv:math/0606241](https://arxiv.org/abs/math/0606241).
- [15] Yu. I. Manin, Noncommutative geometry and quantum theta functions, *Lecture Notes*, Independent Moscow Univ., May 29, 2000. (Russian)

- [16] Yu. I. Manin, Real multiplication and noncommutative geometry, in “Legacy of Niels Hendrik Abel”, 685-727, Springer, 2004.
- [17] H. Masur, Interval exchange transformations and measured foliations, *Ann. of Math. (2)* 115 (1982), no. 1, 169–200.
- [18] O. Perron, Grundlagen für eine Theorie des Jacobischen Kettenbruchalgorithmus, *Math. Annalen* 64 (1907), 1-76.
- [19] A. Polishchuk and A. Schwarz, Categories of holomorphic vector bundles on noncommutative two-tori, *Commun. Math. Phys.* 236 (2003), 135-159.
- [20] Y. Soibelman and V. Vologodsky, Noncommutative compactifications and elliptic curves, *Int. Math. Res. Not.* (2003), 1549-1569.
- [21] K. Strebel, *Quadratic Differentials*, Springer, 1984.
- [22] W. P. Thurston, On the geometry and dynamics of diffeomorphisms of surfaces, *Bull. Amer. Math. Soc.* 19 (1988), 417-431.
- [23] W. A. Veech, Gauss measures for transformations on the space of interval exchange maps, *Ann. of Math. (2)* 115 (1982), no. 1, 201-242.

THE FIELDS INSTITUTE FOR MATHEMATICAL SCIENCES, TORONTO, ON,
CANADA, E-MAIL: igor.v.nikolaev@gmail.com

Current address: 101-315 Holmwood Ave., Ottawa, ON, Canada, K1S 2R2